## **3 EASY STEPS FOR SUPERIOR HOLDING POWER**

Clean and dry the area where Mastisol will be applied. Place an individual vial in the blister-pak on a hard surface (label facing down). Gently apply pressure to the vial through the blister-pak to crush the inner ampule, thereby releasing the Mastisol. Do not crush the ampule more than once.

Remove the individual vial using an aseptic technique. Peel back the lid and deliver the Mastisol medical adhesive vial.

Carefully apply Mastisol by gently squeezing the vial. Apply Mastisol to the area that will be covered by the tape or dressing. Be careful not to come into contact with wound tissue. Allow Mastisol to dry for 20-30 seconds. Using an established technique place the tape or dressing in contact with Mastisol.

Warnings: Contains:

For external use only. Gurn Mastic, Styrax Liquid, SD Alcohol 23A, and Methyl Salicylate

Flammable. Keep out of the reach of Children. Caution:









2 PEEL



3 APPLY

## **COMMON MASTISOL APPLICATION AREAS**



WOUND CLOSURE STRIPS



OSTOMY APPLIANCE



DRESSING



ENDOTRACHEAL TUBE





When you need it on firmly, securely, easily, and safely.



Mastisol is a liquid medical adhesive used to ensure the integrity of dressings, tapes, and certain devices over an extended period of time.