



## **REF CA-42854-P1A**

## Contents:

- 1: Quad-Lumen Indwelling Catheter: 8.5 Fr. x 16 cm Pressure Injectable Radiopague Polyurethane with Blue FlexTip®, ARROWgard Blue PLUS® Antimicrobial Surface Treatment<sup>1</sup>, Extension Line Clamps
- 1: Spring-Wire Guide, Marked: .032" (0.81 mm) dia. x 23-5/8" (60 cm) (Straight Soft Tip on One End - "J" Tip on Other) with Arrow Advancer
- 1: Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer Needle
- 1: Pressure Transduction Probe
- 1: Introducer Needle: Echogenic 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction Syringe
- 1: Injection Needle: Eclipse<sup>™2</sup> 22 Ga. x 1-1/2" (3.81 cm) RW and 5 mL Luer-Slip Syringe
- 1: Injection Needle: Eclipse<sup>™2</sup> 25 Ga. x 1" (2.54 cm) RW and 3 mL Luer-Lock Syringe
- 1: Syringe: 10 mL Luer-Lock

- 1: Tissue Dilator: 10 Fr. (3.5 mm) x 10.2 cm
- 1: SecondSite<sup>™</sup> Adjustable Hub: Fastener
- 1: SecondSite<sup>™</sup> Adjustable Hub: Catheter Clamp
- 1: SharpsAway<sup>®</sup> Disposal Cup
- 1: SharpsAway II<sup>™</sup> Locking Disposal Cup
- 1: Drape: 41" x 55" with 4" fenestration
- 1: Needle Holder
- 1: Safety Scalpel: #11
- 1: Flow Rate Information Card with Injection Log
- 1: Checklist/CLIP Sheet
- 1: Sterile Procedure Sign
- 2: Gauze Pad: 2" x 2" (5 cm x 5 cm)
- 5: Gauze Pad: 4" x 4" (10 cm x 10 cm)
- 1: Dressing: Tegaderm<sup>®3</sup> 10 cm x 12 cm
- 1: Suture: 3-0 Silk with Curved Needle

Contraindications: The ARROWgard Blue Plus antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine and/or sulfa drugs (refer to product instructions for references).

1: HemoHopper<sup>®</sup> Fluid Receptacle

<sup>1</sup>Licensed under US Patent Nos. 6,706,024; 6,872,195 and 7,329,412. <sup>2</sup>A trademark of Becton, Dickinson and Company. <sup>3</sup>A registered trademark of 3M Company.

í			14 Ga.	
	Fr. mm	18 Ga.		18 Ga.
			16 Ga.	

Lumen	Priming Volume* (mL)	Flow Rate† (mL/hr)	MAX Pressure Injection Flow Rate‡ (mL/sec)
Distal (16 Ga.)	0.38	3350	5
Medial 1 (14 Ga.)	0.45	6000	10
Medial 2 (18 Ga.)	0.35	1750	not rated
Proximal (18 Ga.)	0.38	1950	not rated

\* Priming volumes are approximate and are done without injection caps. **Catheter Cross Section** + Based on 100 cm head height.

<sup>‡</sup> Pressure injection flow rates are determined using media of 11.8 centipoise (cP) viscosity, with 60" administration set tubing.

Not made with natural rubber latex

Store below 25°C (77°F). Avoid excessive heat above 40°C (104°F).

A-42854-300A (04/18)



	14 Ga.	
18 Ga.	_	18 Ga.
$\sim$		
	16 Ga.	

		14 Ga.		
r. m	18 Ga.		18 Ga.	
		16Ga		

	14Ga.		Lumen
18 Ga.		18 Ga.	Distal (1 Modial 1

	Priming	Flow

8.5<sup>Fr.</sup> 4<sup>Lumen</sup> 16<sup>cm</sup><sub>catheter</sub> .032<sup>inch dia.</sup>