





Pressure Injectable Two-Lumen CVC Kit



REF CA-46702-P1A

7 Fr. 2 Lumen 20 cm catheter length .032 spring-wire guide

Contents:

- 1: Two-Lumen Indwelling Catheter: 7 Fr. x 20 cm Pressure Injectable Radiopague Polyurethane with Blue FlexTip®, ARROWgard Blue PLUS® Antimicrobial Surface Treatment¹, Extension Line Clamps
- 1: Spring-Wire Guide, Marked: .032" (0.81 mm) dia. x 23-5/8" (60 cm) (Straight Soft Tip on One End - "J" Tip on Other) with Arrow Advancer
- 1: Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer Needle
- 1: Pressure Transduction Probe
- 1: Introducer Needle: Echogenic 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction Syringe
- 1: Injection Needle: Eclipse™2 22 Ga. x 1-1/2" (3.81 cm) RW and 5 mL Luer-Slip Syringe

- 1: Injection Needle: Eclipse^{™2} 25 Ga. x 1" (2.54 cm) RW and 3 mL Luer-Lock Syringe
- 1: Syringe: 10 mL Luer-Lock
- 1: Tissue Dilator: 8.5 Fr. (2.8 mm) x 10.2 cm
- 1: SecondSite™ Adjustable Hub: Fastener
- 1: SecondSite™ Adjustable Hub: Catheter Clamp
- 1: SharpsAway® Disposal Cup
- 1: SharpsAway II™ Locking Disposal Cup
- 1: Drape: 41" x 55" with 4" fenestration
- 1: Needle Holder
- 1: Safety Scalpel: #11
- 1: Checklist/CLIP Sheet

- 1: Sterile Procedure Sign
- 1: Flow Rate Information Card with Injection Log
- 2: Gauze Pad: 2" x 2" (5 cm x 5 cm)
- 5: Gauze Pad: 4" x 4" (10 cm x 10 cm)
- 1: Dressing: Tegaderm®3 10 cm x 12 cm
- 1: Suture: 3-0 Silk with Curved Needle
- 1: HemoHopper® Fluid Receptacle

¹Licensed under US Patent Nos. 6,706,024; 6,872,195 and 7,329,412.

²A trademark of Becton, Dickinson and Company.


³A registered trademark of 3M Company.

Contraindications: The ARROWgard Blue Plus antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine and/or sulfa drugs (refer to product instructions for references).





Store below 25°C (77°F). Avoid excessive heat above 40°C (104°F).



Lumen	Priming Volume* (mL)	Flow Rate† (mL/hr)	MAX Pressure Injection Flow Rate‡ (mL/sec)
Distal (16 Ga.)	0.42	3150	5
Proximal (16 Ga.)	0.45	2800	10

- * Priming volumes are approximate and are done without accessories.
- † Based on 100 cm head height.
- ‡ Pressure injection flow rates are determined using media of 11.8 centipoise (cP) viscosity, with 60" administration set tubing.

A-46702-300A (03/18)