

Pressure Injectable Arrowg+ard Blue Plus® Three-Lumen CVC

REF CDC-42703-P1A

Contents:

- 1: Arrowg+ard Blue Plus®1 Three-Lumen Catheter: 7 Fr. (2.5 mm OD) x 16 cm, Pressure Injectable
- 1: Spring-Wire Guide, Marked: .032" (0.81 mm) dia. x 17-3/4" (45 cm) (Straight Soft Tip on One End - "J" Tip on Other) with Arrow Advancer
- 1: Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer Needle
- 1: Injection Needle: SafetyGlide^{™2} 25 Ga. x 1" (2.54 cm)
- 1: Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm)
- 1: Pressure Transduction Probe
- 1: Introducer Needle: Echogenic 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction Syringe

Contraindications: The Arrowg+ard Blue Plus antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine and/or sulfa drugs (refer to product instructions for references). Arrow International provides the enclosed medication label(s) for your convenience. Please ensure that these labels are

- 1: Injection Needle: SafetyGlide^{™2} 22 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip Syringe
- 1: Syringe: 3 mL Luer-Lock
- 1: Tissue Dilator: 8.5 Fr. (2.8 mm) x 10.2 cm

- 1: 5 mL 1% Lidocaine HCI Solution and Alcohol Prep
- 1: 3 mL Applicator 2% CHG and 70% IPA ChloraPrep® One-Step Solution with Hi-Lite Orange[™] Tint

7^{Fr.} 3^{Lumen} 16^{cm} 13^{cm} 13^{cm} 16^{cm} 3^{cm} 3^{cm}

- 1: SecondSite[™] Adjustable Hub: Fastener
- 1: SecondSite[™] Adjustable Hub: Catheter Clamp
- 1: SharpsAway[®] II Locking Disposal Cup
- 1: SharpsAway[®] Disposal Cup
- 1: Maximal Barrier Drape[™] with 4" fenestration
- 1: Needle Holder
- 1: Safety Scalpel: #11
- 1: Flow Rate Information Card with Injection Log
- 1: Sterile Procedure Sign
- 1: Medication Label: 1% Lidocaine
- 2: Gauze Pad: 2" x 2" (5 cm x 5 cm)
- 5: Gauze Pad: 4" x 4" (10 cm x 10 cm)

- 1: Surgical Apparel: Impervious Gown
- 1: Surgical Apparel: Bouffant Cap
- 1: Surgical Apparel: Mask with Eye Shield
- 1: Suture: 3-0 Silk with Curved Needle

¹Licensed under US Patent No. 7,329,412. ²A trademark of Becton, Dickinson and Company, ³A registered trademark of CareFusion or one of its subsidiaries.

Lumen	Priming Volume* (mL)	Flow Rate† (mL/hr)	MAX Pressure Injection Flow Rate‡ (mL/sec)
Distal (16 Ga.)	0.38	3000	10
Medial (18 Ga.)	0.37	1400	5
Proximal (18 Ga.)	0.39	1600	5

+ Based on 100 cm head height.

[‡] Pressure injection flow rates are determined using media of 11.8 centipoise (cP) viscosity, with 152 cm administration set tubing.

Not made with



applied to the correct syringe and corresponding medication.

natural rubber latex 20-25°C (68-77°F).



C-42703-300A (2020-12)

