







Arrowg+ard Blue® Two-Lumen Hemodialysis **Catheterization Kit for High Volume Infusions**

16 cm length .035 inch dia. spring-w guide

Contents:

- 1: Two-Lumen Indwelling Catheter: 12 Fr. x 16 cm Radiopaque Polyurethane with Blue FlexTip®, Arrowg+ard® Antimicrobial Surface Treatment¹, Extension Line Clamps
- Spring-Wire Guide, Marked: .035" (0.89 mm) dia. x 23-5/8" (60 cm) (Straight Soft Tip on One
- End "L" Tip on Other) with Arrow Advancer Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer Needle
- Injection Needle: 25 Ga. x 1" (2.54 cm)
- Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm)
- Pressure Transduction Probe
- Introducer Needle: 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow $^{\circ}$ Raulerson Spring-Wire Introduction Syringe
- Injection Needle: 22 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip Syringe
- Syringe: 3 mL Luer-Lock
- Tissue Dilator: 12 Fr. (4.0 mm) x 14 cm
- 5 mL 1% Lidocaine HCl Solution and Alcohol Prep
- 3 mL Applicator 2% CHG and 70% IPA ChloraPrep®2 One-Step Solution with Hi-Lite Orange™ Tint (non-sterile solution) Dust Cap: Non-Vented
- SharpsAway II™ Locking Disposal Cup
- SharpsAway® Disposal Cup Drape: 41" x 55" with 4" fenestration
- Needle Holder
- Safety Scalpel: #11 Gauze Pad: 2" x 2" (5 cm x 5 cm)
- Gauze Pad: 4" x 4" (10 cm x 10 cm)
- Suture: 3-0 Silk with Curved Needle

¹Licensed under US Patent No. 6,706,024.


Rx only. Warning: Read all package insert warnings, cautions, and instructions prior to use. Failure to do so may result in severe patient injury or death.

⚠ California Prop. 65

Warning: Cancer and Reproductive Harm www.P65Warnings.ca.gov.

Sterilized using ethylene oxide unless otherwise indicated in the contents list. Fluid path components are non-pyrogenic.

Contraindications: The Arrow large-bore two-lumen catheter is not designed for long-term hemodialysis or for use in patients with thrombosed vessels. The Arrowg+ard Blue antimicrobial surface catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine, and/or sulfa drugs (refer to product instructions for references).



| Lumen | Priming Volume* (mL) | Flow Rate† (mL/min) | Venous Pressure Setting (mmHg) |
|-------------------|----------------------------|---------------------------|--------------------------------------|
| Distal (12 Ga.) | 1.40 | 220 | 100 |
| | | 370 | 200 |
| Proximal (12 Ga.) | 1.30 | 220 | 100 |
| | | 370 | 200 |

* Priming volumes are approximate and are done without accessories. † Blood flow rates are approximate and were determined by varying the venous pressure setting from 100 to 200 mmHg in an in vitro experiment using blood analog solution.





















Arrow International LLC Subsidiary of Teleflex Incorporated 3015 Carrington Mill Blvd. Morrisville, NC 27560 USA



IK-22122-105A, Rev. 01 (2020-12)

REF AK-22122-CDC







