

Arrowg+ard Blue Advance[™] PICC pre-loaded with Arrow[®] VPS Precision[™] Stylet

REE CDC-45552-VPS2

Contents:

- 1: Two-Lumen TaperFree[®] Catheter with Arrowg+ard Blue Advance™ Antimicrobial/Antithrombogenic Protection¹: 5.5 Fr. x 55 cm Pressure Injectable with Blue FlexTip[®] Feature, Arrow[®] VPS Precision[™] Stylet, Tuohy-Borst Adapter±, Marking Device, and Console Pointers
- 1: GlideThru[™] Peel-Away Sheath: 5.5 Fr. x 2-3/4" (7 cm) Radiopague over 5.5 Fr. Dilator
- 1: Spring-Wire Guide, Nitinol, Marked: .018" (0.46 mm) dia, x 17-3/4" (45 cm) (Straight Soft Tip on One End - Straight Stiff Tip on Other) with Arrow Advancer
- 1: Injection Needle: SafetvGlide^{™2} 25 Ga. x 5/8" (1.60 cm) RW
- 1: Safety Introducer Needle: Echogenic 21 Ga. x 2-3/4" (7 cm) TW
- 1: Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm)
- 1: Introducer Needle: Echogenic 21 Ga. x 2-3/4" (7 cm) TW
- 1: Syringe: 10 mL Luer-Lock
- 1: Syringe: 3 mL Luer-Lock
- 1: Skin Protectant Prep Pad (non-sterile solution)
- 4: 10 mL Luer-Lock Pre-Filled Saline Syringe
- 1: 5 mL 1% Lidocaine HCl Solution
- 1: 3 mL Applicator 2% CHG and 70% IPA ChloraPrep® One-Step Solution (non-sterile solution)

with pregnant women, nursing mothers, babies and infants.

applied to the correct syringe and corresponding medication.

1: 3 mL Applicator 2% CHG and 70% IPA ChloraPrep® One-Step Solution with Hi-Lite Orange[™] Tint (non-sterile solution)

Sterilized using ethylene oxide unless otherwise indicated in the contents list.

thrombosis in the intended vessel or along the catheterized vessel pathway.

DEHP

[±] Precaution:

- 2: Dust Cap: Non-Vented
- 1: SecondSite[™] Adjustable Hub: Fastener
- 1: SecondSite[™] Adjustable Hub: Catheter Clamp
- 1: SharpsAway® Disposal Cup
- 1: SharpsAway II[™] Locking Disposal Cup
- 1: Catheter Trimmer
- 1: Towel: 24" x 36" (61 cm x 91 cm)
- 1: Drape: 60" x 76" (152 cm x 193 cm)
- 1: Drape: 36" x 41" (91 cm x 104 cm) with 3" x 5" (8 cm x 13 cm) fenestration

5.5^{Fr.} 2^{Lumen} 55^{cm} 018^{inch dia.}

- 1: Safety Scalpel: #11
- 1: Sterile Procedure Sign
- 1: Patient ID Card
- 1: Chart Sticker

Towel

- 1: Patient Information Booklet
- 1. Medication Label: 1% Lidocaine
- 1: Checklist/CLIP Sheet
- 1: Flow Rate Information Card with Injection Log
- 2: Paper Tape Measure
- 1: Tourniquet
- 1: Foam Electrodes (3 per pack)
- 10: Gauze Pad: 2" x 2" (5 cm x 5 cm)

- Erg[®]Pack
- 10: Gauze Pad: 4" x 4" (10 cm x 10 cm)
- 1: Dressing: Tegaderm®4 15.5 cm x 10 cm
- 1: Surgical Apparel: Impervious Gown
- 1: Dressing: BIOPATCH®5
- 1: Dressing: STATLOCK®6 Catheter Stabilization Device
- Surgical Apparel: Mask with Eye Shield 1.
- 1: Surgical Apparel: Bouffant Cap
- 1: Surgical Apparel: Mask
- 1: Tape: Steri-Strip^{®4}

Lumen

Distal (18 Ga.)

Proximal (18 Ga.)

1: HemoHopper® Fluid Receptacle

¹Licensed under US Patent Nos. 6,706,024, 6,872,195 and 7,329,412. ²A trademark of Becton, Dickinson and Company. ³A registered trademark of CareFusion or one of its subsidiaries. ⁴A registered trademark of 3M Company. ⁵A registered trademark of Johnson & Johnson Corporation. 6A registered trademark of C. R. Bard, Inc.

Priming

Volume³

(mL)

0.49

0.50

hypersensitivity to chlorhexidine • In presence of device related infections • In presence of previous or current 5.5 Fr. Warning: The results of certain animal experiments have shown phthalates to be potentially toxic to reproduction. 1.9 mn

370 * Priming volumes are approximate and are done without accessories. ⁺ Flow rates were determined using room temperature water, 100 cm head height and represent approximate flow capabilities.

Flow

Rate

(mL/hr)

360

Pump Flow

Rate[‡]

(mL/hr)

3020

2970

tion

- * Pump flow rates are determined at maximum pump pressure of 10 psig and represent approximate flow capabilities.
- ** Pressure injection flow rates are determined at the injector pressure setting of 300 psi maximum using media of 11.8 centipoise viscosity, with 152 cm pressure tubing.

IV-45552-330B (11/18)

MAX Pressure

Injection

Flow Rate**

(mL/sec)

5

5

Not made with natural rubber latex



Store between 20-25°C (68-77°F)

Contraindications: Clinical assessment of the patient must be completed to ensure no contraindications exist. The

Arrow PICC with Arrowg+ard Blue Advance Protection is contraindicated in the following areas: • Patients with known

Proceeding from the present state of scientific knowledge, risks for male premature infants cannot be excluded in the

case of long-term exposure or application. Medical products containing phthalates should be used only temporarily

Arrow International provides the enclosed medication label(s) for your convenience. Please ensure that these labels are

18Ga 1000

,		o Ga.
	Catheter Cross S	Sect