



16 cm catheter length

■032 inch dia. spring-wire guide



Arrowg+ard Blue Plus® Two-Lumen CVC

Contents:

- 1: Arrowg+ard Blue Plus®1 Two-Lumen Catheter: 8 Fr. (2.8 mm OD) x 16
- Spring-Wire Guide, Marked: .032" (0.81 mm) dia. x 17-3/4" (45 cm) (Straight Soft Tip on One End - "J" Tip on Other) with Arrow Advancer
- Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer Needle
- 1: Injection Needle: SafetyGlide™2 25 Ga. x 1" (2.54 cm)
- 1: Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm)
- 1: Pressure Transduction Probe
- Introducer Needle: Echogenic 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction Syringe
- Injection Needle: SafetyGlide™2 22 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip Syringe
- 1: Syringe: 3 mL Luer-Lock
- 1: Tissue Dilator: 9 Fr. (3.0 mm) x 10.2 cm
- 1: 5 mL 1% Lidocaine HCI Solution and Alcohol Prep
- 3 mL Applicator 2% CHG and 70% IPA ChloraPrep®3 One-Step Solution with Hi-Lite Orange™ Tint
- SecondSite™ Adjustable Hub: Fastener
- SecondSite™ Adjustable Hub: Catheter Clamp
- SharpsAway® Disposal Cup
- SharpsAway® II Locking Disposal Cup
- Maximal Barrier Drape™ with 4" fenestration
- Needle Holder
- Safety Scalpel: #11
- Checklist/CLIP Sheet
- Sterile Procedure Sign

- 1: Medication Label: 1% Lidocaine
- 2: Gauze Pad: 2" x 2" (5 cm x 5 cm)
- Gauze Pad: 4" x 4" (10 cm x 10 cm)
- Surgical Apparel: Impervious Gown
- 1: Surgical Apparel: Bouffant Cap
- Surgical Apparel: Mask with Eye Shield
- Suture: 3-0 Silk with Curved Needle ¹Licensed under US Patent No. 7,329,412.
- 1: HemoHopper® Fluid Receptacle
- ²A trademark of Becton, Dickinson and Company,
- 3A registered trademark of CareFusion or one of its subsidiaries.

Not made with natural rubber latex.
Store between 20 – 25°C (68 – 77°F).

Arrow International provides the enclosed medication label(s) for your convenience. Please ensure that these labels are applied to the correct syringe and corresponding medication.

Contraindications: The Arrowg+ard Blue Plus antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine and/or sulfa drugs.

Lumen	Priming Volume* (mL)	Gravity Flow Rate† (mL/hr)	
Distal (14 Ga.)	0.68	6350	
Proximal (14 Ga.)	0.70	4980	

^{*} Priming volumes are approximate and are done without accessories.



LBL052799 R00 (2021-10)

[†] Flow rate values are approximate and are determined using deionized water at 100 cm head