







(IPNXXXXXX)



Pressure Injectable Arrowg+ard Blue Plus® Two-Lumen CVC



Contents:

- Arrowg+ard Blue Plus®1 Two-Lumen Catheter: 8 Fr. (2.8 mm OD) x 20 cm, Pressure Injectable Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXX
- Spring-Wire Guide, Marked: .032" (0.81 mm) dia. x 23-5/8" (60 cm) (Straight Soft Tip on One End - "J" Tip on Other) with Arrow Advancer Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXX
- Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer Needle Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXX
- Injection Needle: SafetyGlide™2 25 Ga. x 1" (2.54 cm) Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXXX
- Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm) Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXXX
- Pressure Transduction Probe Origin: XX XXXX XX XXX XXXXXXX
- Introducer Needle: Echogenic 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction Syringe Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXXX
- Injection Needle: SafetyGlide™2 23 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip Syringe Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXX
- Tissue Dilator: 9 Fr. (3.0 mm) x 10.2 cm Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXXX
- 10 mL Luer-Lock Pre-Filled Saline Syringe Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXXX
- 5 mL 1% Lidocaine HCl Solution and Alcohol Prep Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXXX
- 3 mL Applicator 2% CHG and 70% IPA ChloraPrep®3 One-Step Solution with Hi-Lite Orange™ Tint Origin: XX XXXX XX XXXXXXXX XXXXXXXXXXX

- SecondSite™ Adjustable Hub: Fastener Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXXX
- SecondSite™ Adjustable Hub: Catheter Clamp Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXXX
- SharpsAway® II Locking Disposal Cup Origin: XX XXXX XX XXX XXXXXXX
- SharpsAway® Disposal Cup Origin: XX XXXX XX XXX XXXXXXX
- Maximal Barrier Drape™ with 4" fenestration Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXXX
- Towel Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXX
- Needle Holder Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXX
- Safety Scalpel: #11 Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXX
- Flow Rate Information Card with Injection Log Origin: XX XXXX XX XXX
- XXXXXXX XXXXXXXXXX Sterile Procedure Sign Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXX
- Medication Label: 1% Lidocaine Origin: XX XXXX XX XXX XXXXXXX
- XXXXXXXXXX Gauze Pad: 2" x 2" (5 cm x 5 cm) Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXXX
- Gauze Pad: 4" x 4" (10 cm x 10 cm) Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXXX
- Dressing: Tegaderm®4 10 cm x 12 cm Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXX
- Surgical Apparel: Impervious Gown Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXXX

- Dressing: BIOPATCH®5 Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXX
- Surgical Apparel: Bouffant Cap Origin: XX XXXX XX XXX XXXXXXX
- Surgical Apparel: Mask with Eye Shield Origin: XX XXXX XX XXX XXXXXXX XXXXXXXXX
- Suture: 3-0 Silk with Curved Needle Origin: XX XXXX XX XXX XXXXXXX
- XXXXXXXXX Tubing±: 20" (50 cm) Extension Origin: XX XXXX XX XXX XXXXXXX
- XXXXXXXXX HemoHopper® Fluid Receptacle Origin: XX XXXX XX XXX XXXXXXX
- XXXXXXXXXX
- ¹Licensed under US Patent No. 7,329,412.
- ²A trademark of Becton, Dickinson and Company.
- 3A registered trademark of CareFusion or one of its subsidiaries.
- ⁴A registered trademark of 3M Company.
- 5A registered trademark of Johnson & Johnson Corporation


±Precaution: Contains Phthalate: DEHP.

Warning: The results of certain animal experiments have shown phthalates to be potentially toxic to reproduction. Proceeding from the present state of scientific knowledge, risks for male premature infants cannot be excluded in the case of long-term exposure or application. Medical products containing phthalates should be used only temporarily with pregnant women, nursing mothers, babies and infants. Not made with natural rubber latex.

Store between 20 - 25°C (68 - 77°F)

Arrow International provides the enclosed medication label(s) for your convenience. Please ensure that these labels are applied to the correct syringe and

Contraindications: The Arrowg+ard Blue Plus antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine and/or sulfa drugs.



- * Priming volumes are approximate and are done without accessories.
- † Flow rate values are approximate and are determined using deionized water at 100 cm head
- ** Pressure injection flow rates are determined using media of 11.8 centipoise (cP) viscosity, with 152 cm administration set tubing.











LBL052967 R02 (2025-03)