



REF CDC-45802-XP1A (IPNXXXXXX)

8 Fr. 2 Lumen 20 cm catheter length .032 inch dia. spring-wire guide



Pressure Injectable Arrowg+ard Blue Plus® Two-Lumen CVC



Contents:

- 1: Arrowg+ard Blue Plus®¹ Two-Lumen Catheter: 8 Fr. (2.8 mm OD) x 20 cm, Pressure Injectable Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Spring-Wire Guide, Marked: .032" (0.81 mm) dia. x 23-5/8" (60 cm) (Straight Soft Tip on One End - "J" Tip on Other) with Arrow Advancer Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer Needle Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Injection Needle: SafetyGlide™² 25 Ga. x 1" (2.54 cm) Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm) Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Pressure Transduction Probe Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Introducer Needle: Echogenic 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction Syringe Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Injection Needle: SafetyGlide™² 23 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip Syringe Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Syringe: 3 mL Luer-Lock Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Tissue Dilator: 9 Fr. (3.0 mm) x 10.2 cm Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 2: 10 mL Luer-Lock Pre-Filled Saline Syringe Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: 5 mL 1% Lidocaine HCl Solution and Alcohol Prep Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: 3 mL Applicator 2% CHG and 70% IPA Chloraprep®³ One-Step Solution with Hi-Lite Orange™ Tint Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: SecondSite™ Adjustable Hub: Fastener Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: SecondSite™ Adjustable Hub: Catheter Clamp Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: SharpsAway® II Locking Disposal Cup Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: SharpsAway® Disposal Cup Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Maximal Barrier Drape™ with 4" fenestration Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Towel Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Needle Holder Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Safety Scalpel: #11 Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Checklist/CLIP Sheet Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Flow Rate Information Card with Injection Log Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Sterile Procedure Sign Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Medication Label: 1% Lidocaine Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 2: Gauze Pad: 2" x 2" (5 cm x 5 cm) Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 5: Gauze Pad: 4" x 4" (10 cm x 10 cm) Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Dressing: Tegaderm®⁴ 10 cm x 12 cm Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Surgical Apparel: Impervious Gown Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Dressing: BIOPATCH®⁵ Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Surgical Apparel: Bouffant Cap Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Surgical Apparel: Mask with Eye Shield Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Suture: 3-0 Silk with Curved Needle Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: Tubing±: 20" (50 cm) Extension Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX
- 1: HemoHopper® Fluid Receptacle Origin: XX XXXX XX XXX XXXXXXXXXXXXXXXXXXXX



Lumen	Priming Volume* (mL)	Gravity Flow Rate† (mL/hr)	MAX Pressure Injection Flow Rate** (mL/sec)
Distal (14 Ga.)	0.73	5354	10
Proximal (14 Ga.)	0.75	3984	10

* Priming volumes are approximate and are done without accessories.
 † Flow rate values are approximate and are determined using deionized water at 100 cm head height.
 ** Pressure injection flow rates are determined using media of 11.8 centipoise (cP) viscosity, with 152 cm administration set tubing.

⚠Precaution: Contains Phthalate: DEHP.
Warning: The results of certain animal experiments have shown phthalates to be potentially toxic to reproduction. Proceeding from the present state of scientific knowledge, risks for male premature infants cannot be excluded in the case of long-term exposure or application. Medical products containing phthalates should be used only temporarily with pregnant women, nursing mothers, babies and infants.
Not made with natural rubber latex.
 Store between 20 – 25°C (68 – 77°F).
 Arrow International provides the enclosed medication label(s) for your convenience. Please ensure that these labels are applied to the correct syringe and corresponding medication.
Contraindications: The Arrowg+ard Blue Plus antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine and/or sulfa drugs.

