







(IPNXXXXXX)

Pressure Injectable Arrowg+ard Blue Plus® Three-Lumen CVC



Contents:

- 1: Arrowg+ard Blue Plus®1 Three-Lumen Catheter: 7 Fr. (2.5 mm OD) x 20 cm, Pressure Injectable
- Spring-Wire Guide, Marked: .032" (0.81 mm) dia. x 23-5/8" (60 cm) (Straight Soft Tip on One End - "J" Tip on Other) with Arrow Advancer
- Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer
- Pressure Transduction Probe
- Introducer Needle: Echogenic 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction Syringe
- Injection Needle: Eclipse™2 23 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip
- Injection Needle: Eclipse™2 25 Ga. x 1" (2.54 cm) and 3 mL Luer-Lock Syringe
- Syringe: 10 mL Luer-Lock
- Tissue Dilator: 8.5 Fr. (2.8 mm) x 10.2 cm
- SecondSite™ Adjustable Hub: Fastener
- SecondSite™ Adjustable Hub: Catheter Clamp
- SharpsAway® II Locking Disposal Cup
- SharpsAway® Disposal Cup
- Drape: 41" x 55" with 4" fenestration
- Needle Holder
- Safety Scalpel: #11
- Checklist/CLIP Sheet
- Sterile Procedure Sign
- Gauze Pad: 2" x 2" (5 cm x 5 cm)
- Gauze Pad: 4" x 4" (10 cm x 10 cm)
- Dressing: Tegaderm®3 10 cm x 12 cm
- 1: Suture: 3-0 Silk with Curved Needle

- 1: HemoHopper® Fluid Receptacle
- ¹Licensed under US Patent No. 7,329,412.
- ²A trademark of Becton, Dickinson and Company
- ³A registered trademark of 3M Company

Not	made	with	natural	rubber	latex.

Store below 25°C (77°F). Avoid excessive heat above 40°C (104°F)

Contraindications: The Arrowg+ard Blue Plus antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine and/or sulfa drugs.

Lumen	Priming Volume* (mL)	Gravity Flow Rate† (mL/hr)	MAX Pressure Injection Flow Rate** (mL/sec)
Distal (16 Ga.)	0.39	2283	10
Medial (18 Ga.)	0.39	913	5
Proximal (18 Ga.)	0.40	996	5

^{*} Priming volumes are approximate and are done without accessories.



LBL053589 R02 (2024-05)

[†] Flow rate values are approximate and are determined using deionized water at 100 cm head

^{**} Pressure injection flow rates are determined using media of 11.8 centipoise (cP) viscosity, with 152 cm administration set tubing.