

1:

Needle Holder

Pressure Injectable Arrowg+ard Blue Plus® Two-Lumen CVC

SharpsAway[®] II Locking Disposal Cup Drape: 41" x 55" with 4" fenestration

Medication Label: 1% Lidocaine

¹Licensed under US Patent No. 7.329,412. ²A trademark of Becton, Dickinson and Company

Safety Scalpel: #11 Flow Rate Information Card with Injection Log Sterile Procedure Sign Central Line Insertion Checklist

³A registered trademark of CareFusion or one of its subsidiaries.

Contents

- 1: Arrowg+ard Blue Plus®1 Two-Lumen Catheter: 8 Fr. (2.8 mm OD) x 20 cm,
- Pressure Injectable Spring-Wire Guide, Marked: .032" (0.81 mm) dia, x 23-5/8" (60 cm) (Straight
- 1: Soft Tip on One End - "J" Tip on Other) with Arrow Advancer Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer 1:
- Needle
- Injection Needle: SafetyGlide™2 25 Ga. x 1" (2.54 cm) Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm)
- 1:
- 1:
- Pressure Transduction Probe
 1.
 Inteducation Laber. 17% Eldocatine

 Introducer Needle: Echogenic 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL
 2.
 Gauze Pad: 2" x 2" (5 cm x 5 cm)

 Arrow® Raulerson Spring-Wire Introduction Syringe
 1.
 Suture: 3-0 Silk with Curved Needle

 Injection Needle: SafetyGlide^{™2} 22 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip
 1.
 Tubing±: 20" (50 cm) Extension

 1:
- Syringe Syringe: 3 mL Luer-Lock
- Tissue Dilator: 9 Fr. (3.0 mm) x 10.2 cm
- Insue Dilator: 9 Fr. (3.0 mm) x 10.2 cm
 5 mL 1% Lidocaine HCI Solution and Alcohol Prep
 3 mL Applicator 2% CHG and 70% IPA ChloraPrep®3 One-Step Solution with Hi-Lite Orange™ Tint
 SecondSite™ Adjustable Hub: Fastener
 SecondSite™ Adjustable Hub: Catheter Clamp
 SharpsAway® Disposal Cup

Rx only

Warning: Read all package insert warnings, precautions, and instructions prior to use. Failure to do so may result in severe patient injury or death.

California Prop. 65

More and Reproductive Harm. www.P65Warnings.ca.gov Precaution: Contains Phthalate: DEHP.

Warning: The results of certain animal experiments have shown phthalates to be potentially toxic to reproduction. Proceeding from the present state of scientific knowledge, risks for male premature infants cannot be excluded in the case of long-term exposure or application. Medical products containing phthalates should be used only temporarily with pregnant women, nursing mothers, babies and infants.

Store between 20 – 25°C (68 – 77°F).

Sterilized using ethylene oxide unless otherwise indicated in the contents list.

Fluid path components are non-pyrogenic. Arrow International provides the enclosed medication label(s) for your convenience. Please ensure that these labels are applied to the correct syringe and corresponding medication.

Contraindications: The Arrowg+ard Blue Plus antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine and/or sulfa drugs.

Q	Priming Volume* (mL)	Gravity Flow Rate† (mL/hr)	MAX Pressure Injection Flow Rate** (mL/sec)
Distal (14 Ga.)	0.73	5354	10
Proximal (14 Ga.)	0.75	3984	10

ARROW

+ Flow rate values are approximate and are determined using deionized water at 100 cm head height.

** Pressure injection flow rates are determined using media of 11.8 centipoise (cP) viscosity, with 152 cm administration set tubing.



 \sim

Arrow International LLC Subsidiary of Teleflex Incorporated 3015 Carrington Mill Blvd. Morrisville, NC 27560 USA



LBL053633 R00 (2022-01)

ARROW



 \sum

Pressure Injectable Arrowg+ard Blue Plus® Two-Lumen **CVC**

8^{Fr.}

 2^{Lumen}

20 cm catheter .032 spring-wire cuide