





16 cm catheter length 1032 inch dia. spring-wire guide

Pressure Injectable Three-Lumen CVC

- Three-Lumen Catheter: 7 Fr. (2.5 mm OD) x 16 cm, Pressure Injectable Spring-Wire Guide, Marked: .032" (0.81 mm) dia. x 23-5/8" (60 cm) (Straight Soft Tip on One End "J" Tip on Other) with Arrow Advancer
- Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer

- Needle
 Injection Needle: SafetyGlide™1 25 Ga. x 1" (2.54 cm)
 Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm)
 Pressure Transduction Probe
 Introducer Needle: Echoenic 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL
 Arrow® Raulerson Spring-Wire Introduction Syringe
 Injection Needle: SafetyGlide™1 22 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip
 Syringe
 Syringe: 3 mL Luer-Lock

 1. Flow Kate Information Card with Injection Log
 Medication Label: 1% Lidocaine
 12. Gauze Pad: 2" x 2" (5 cm x 5 cm)
 13. Suture: 3-0 Silk with Curved Needle
 14. Tubing±: 20" (50 cm) Extension
 14. trademark of Becton, Dickinson and Company.
 24. registered trademark of CareFusion or one of its subsidiaries.

- Syringe: 3 mL Luer-Lock Tissue Dilator: 8.5 Fr. (2.8 mm) x 10.2 cm 5 mL 1% Lidocaine HCl Solution and Alcohol Prep
- 5 mL 1% Lidocaine HCI Solution and Alcohol Prep
 3 mL Applicator 2% CHG and 70% IPA ChloraPrep®2 One-Step Solution with Hi-Lite Orange™ Tint
 SecondSite™ Adjustable Hub: Fastener
 SecondSite™ Adjustable Hub: Catheter Clamp
 SharpsAway® Disposal Cup
 SharpsAway® II Locking Disposal Cup

Flow Rate Information Card with Injection Log Medication Label: 1% Lidocaine
Gauze Pad: 2" x 2" (5 cm x 5 cm)
Gauze Pad: 4" x 4" (10 cm x 10 cm)

Drape: 41" x 55" with 4" fenestration Needle Holder

Safety Scalpel: #11

Rx only

Warning: Read all package insert warnings, precautions, and instructions prior to use. Failure to do so may result in severe patient injury or death.

California Prop. 65

WARNING: Cancer and Reproductive Harm.
 www.P65Warnings.ca.gov
 Precaution: Contains Phthalate: DEHP.

Warning: The results of certain animal experiments have shown phthalates to be potentially toxic to reproduction. Proceeding from the present state of scientific knowledge, risks for male premature infants cannot be excluded in the case of long-term exposure or application. Medical products containing phthalates should be used only temporarily with pregnant women, nursing mothers, babies and infants.

Not made with natural rubber latex.

Store between 20 - 25°C (68 - 77°F).

Sterilized using ethylene oxide unless otherwise indicated in the contents list.

Fluid path components are non-pyrogenic

Arrow International provides the enclosed medication label(s) for your convenience. Please ensure that these labels are applied to the correct syringe and corresponding medication

Lumen	Priming Volume* (mL)	Gravity Flow Rate† (mL/hr)	MAX Pressure Injection Flow Rate** (mL/sec)
Distal (16 Ga.)	0.4	2947	10
Medial (18 Ga.)	0.4	1552	5
Proximal (18 Ga.)	0.4	1702	5







Arrow International LLC Subsidiary of Teleflex Incorporated 3015 Carrington Mill Blvd. Morrisville, NC 27560 USA



LBL053636 R00 (2021-11)











1032 inch dia. spring-wire quide





[†] Flow rate values are approximate and are determined using deionized water at 100 cm head height.

^{***} Pressure injection flow rates are determined using media of 11.8 centipoise (cP) viscosity, with 152 cm administration set tubing.