



REF AK-25142-CDC

14 Fr. 2 Lumen 20 cm catheter length .038 inch dia. spring-wire guide

Arrowg+ard Blue® Two-Lumen Hemodialysis Catheterization for High Volume Infusions

Contents:

- 1: Two-Lumen Catheter: 14 Fr. (4.72 mm OD) x 20 cm Radiopaque Polyurethane with Blue FlexTip®, Arrowg+ard® Antimicrobial Surface Treatment, Extension Line Clamps
- 1: Spring-Wire Guide, Marked: .038" (0.97 mm) dia. x 27-1/2" (70 cm) (Straight Soft Tip on One End - "J" Tip on Other) with Arrow Advancer
- 1: Injection Needle: 25 Ga. x 1" (2.54 cm)
- 1: Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm)
- 1: Pressure Transduction Probe
- 1: Injection Needle: 22 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip Syringe
- 1: Introducer Needle: 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction Syringe
- 1: Syringe: 3 mL Luer-Lock
- 1: Tissue Dilator: 14 Fr. (4.7 mm) x 14 cm
- 1: Tissue Dilator: 16 Fr. (5.3 mm) x 11.4 cm
- 1: 5 mL 1% Lidocaine HCl Solution and Alcohol Prep
- 1: 3 mL Applicator 2% CHG and 70% IPA Chloraprep®¹ One-Step Solution with Hi-Lite Orange™ Tint
- 2: Dust Cap: Non-Vented
- 1: SharpsAway® II Locking Disposal Cup
- 1: SharpsAway® Disposal Cup
- 1: Drape: 41" x 55" with 4" fenestration
- 1: Needle Holder
- 1: Safety Scalpel: #11
- 2: Gauze Pad: 2" x 2" (5 cm x 5 cm)
- 5: Gauze Pad: 4" x 4" (10 cm x 10 cm)
- 1: Suture: 3-0 Silk with Curved Needle

¹A registered trademark of CareFusion or one of its subsidiaries.

Rx only

Warning: Read all package insert warnings, precautions, and instructions prior to use. Failure to do so may result in severe patient injury or death. California Prop. 65

⚠ **WARNING:** Cancer and Reproductive Harm. www.P65Warnings.ca.gov

Not made with natural rubber latex.

Store between 20 – 25°C (68 – 77°F).

Sterilized using ethylene oxide unless otherwise indicated in the contents list.

Fluid path components are non-pyrogenic.

Contraindications: The Arrow large-bore two-lumen catheter is not designed for long-term (≥ 30 days) hemodialysis or for use in patients with thrombosed vessels.


The Arrowg+ard Blue antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine, and/or sulfa drugs.



Lumen	Priming Volume* (mL)	Flow Rate (mL/min)	Venous Pressure Setting†† (mmHg)
Distal (10 Ga.)	1.7	250	100
		400	170
Proximal (10 Ga.)	1.5	250	120
		400	190

* Priming volumes are approximate and are done without accessories.

†† Venous Pressure settings are approximate and were determined by varying the blood flow rates from 250 to 400 mL/min in an in vitro experiment using blood analog solution.



LOT



Arrow International LLC
Subsidiary of Teleflex Incorporated
3015 Carrington Mill Blvd.
Morrisville, NC 27560 USA

Teleflex®
www.teleflex.com

LBL057086 R00 (2022-02)



REF AK-25142-CDC

14 Fr. 2 Lumen 20 cm catheter length .038 inch dia. spring-wire guide

Arrowg+ard Blue® Two-Lumen Hemodialysis Catheterization for High Volume Infusions