





(IPNXXXXXX)



11.5 cm device length

.035 inch dia. spring-wire guide



## Arrowg+ard Blue® MAC™ Two-Lumen Central Venous Access for use with 7.5 - 8 Fr. **Catheters**



- 1: Two-Lumen Central Venous Access Device: 115 mm Radiopaque Polyurethane with 9 Fr. Distal Lumen, Arrowg+ard® Antimicrobial Surface Treatment1, Integral Hemostasis Valve, Extension Line Clamps
- Spring-Wire Guide, Marked: .035" (0.89 mm) dia. x 17-3/4" (45 cm) (Straight Soft Tip on One End - "J" Tip on Other) with Arrow Advancer
- Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer Needle
- Injection Needle: SafetyGlide™2 25 Ga. x 1" (2.54 cm)
- Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm)
- Pressure Transduction Probe
- Introducer Needle: Echogenic 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction Syringe
- Injection Needle: SafetyGlide™2 23 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip
- Syringe: 3 mL Luer-Lock
- Tissue Dilator
- 5 mL 1% Lidocaine HCl Solution and Alcohol Prep
- 3 mL Applicator 2% CHG and 70% IPA ChloraPrep®3 One-Step Solution with Hi-Lite Orange™ Tint
- Obturator: 8 Fr.
- Cath-Gard® Catheter Contamination Shield: 80 cm with TwistLock™ Adapter

- Dust Cap: Vented
- Dust Cap: Non-Vented
- SharpsAway® II Locking Disposal Cup
- SharpsAway® Disposal Cup
- Maximal Barrier Drape™ with 4" fenestration
- Needle Holder
- Safety Scalpel: #11
- Stopcock: 4-way High-Flow
- Checklist/CLIP Sheet
- Sterile Procedure Sign
- Medication Label: 1% Lidocaine
- Gauze Pad: 2" x 2" (5 cm x 5 cm)
- Gauze Pad: 4" x 4" (10 cm x 10 cm)
- Surgical Apparel: Impervious Gown
- Surgical Apparel: Bouffant Cap
- Surgical Apparel: Mask with Eye Shield
- Suture: 3-0 Silk with Curved Needle Tubing±: 20" (50 cm) Extension
- 1: HemoHopper® Fluid Receptacle

<sup>1</sup>Licensed under US Patent No. 6,706,024.

<sup>2</sup>A trademark of Becton, Dickinson and Company

## ±Precaution: Contains Phthalate: DEHP.

Warning: The results of certain animal experiments have shown phthalates to be potentially toxic to reproduction. Proceeding from the present state of scientific knowledge, risks for male premature infants cannot be excluded in the case of long-term exposure or application. Medical products containing phthalates should be used only temporarily with pregnant women, nursing mothers, babies and infants.

Store between 20 - 25°C (68 - 77°F)

Arrow International provides the enclosed medication label(s) for your convenience. Please ensure that these labels are applied to the correct syringe and

Contraindications: The Arrowg+ard Blue antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine acetate, silver sulfadiazine, and/or sulfa drugs.

Lumen	Priming Volume* (mL)	Gravity Flow Rate† (mL/hr)
Distal (9 Fr.) no catheter Distal (9 Fr.) with 7 Fr. catheter Distal (9 Fr.) with 8 Fr. catheter Proximal (12 Ga.) no catheter	1.7 1.55 1.2 0.55	29310 12440 6720 6710

3A registered trademark of CareFusion or one of its subsidiaries.















LBL058452 R02 (2024-05)

<sup>\*</sup> Priming volumes are approximate and are done without accessories

<sup>†</sup> Flow rate values are approximate and are determined using deionized water at 100 cm head height.