

1.

1:

1:

1:

1.

Transducer Cover: 14 cm x 147 cm

Suture: 3-0 Silk with Curved Needle

1: Tubina±: 20" (50 cm) Extension

HemoHopper[®] Fluid Receptacle

⁴A registered trademark of 3M Company

¹Licensed under US Patent No. 7,329,412.

Surgical Apparel: Mask with Eye Shield Surgical Apparel: Bouffant Cap

²A trademark of Becton, Dickinson and Company

³A registered trademark of CareFusion or one of its subsidiaries.

- Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm) Pressure Transduction Probe
- Introducer Needle: Echogenic 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction Syringe
- Injection Needle: SafetyGlide^{™2} 22 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip Svrinae
- Syringe: 3 mL Luer-Lock
- Tissue Dilator: 9 Fr. (3.0 mm) x 10.2 cm 1.
- 10 mL Luer-Lock Pre-Filled Saline Syringe
- 5 mL 1% Lidocaine HCI Solution and Alcohol Prep
- 3 mL Applicator 2% CHG and 70% IPA ChloraPrep®3 One-Step Solution with 2 Hi-Lite Orange™ Tint
- SecondSite[™] Adjustable Hub: Fastener
- SecondSite[™] Adjustable Hub: Catheter Clamp 1.
- SharpsAway® II Locking Disposal Cup
- SharpsAway® Disposal Cup
- Maximal Barrier Drape™ with 4" fenestration 1.
- Towel 1:
- 1: Needle Holder

±Precaution: Contains Phthalate: DEHP.

Warning: The results of certain animal experiments have shown phthalates to be potentially toxic to reproduction. Proceeding from the present state of scientific knowledge, risks for male premature infants cannot be excluded in the case of long-term exposure or application. Medical products containing phthalates should be used only temporarily with pregnant women, nursing mothers, babies and infants.

Not made with natural rubber latex. Store between 20 - 25°C (68 - 77°F)

Arrow International provides the enclosed medication label(s) for your convenience. Please ensure that these labels are applied to the correct syringe and corresponding medication.

Contraindications: The Arrowg+ard Blue Plus antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine and/or sulfa drugs.



| Q | Priming Volume* (mL) | Gravity Flow Rate† (mL/hr) | MAX Pressure Injection Flow Rate** (mL/sec) |
|-------------------|----------------------------|-------------------------------------|--|
| Distal (14 Ga.) | 0.73 | 5354 | 10 |
| Proximal (14 Ga.) | 0.75 | 3984 | 10 |

| * Priming volumes are | annravimata | and are don | o without | annonarian |
|-----------------------|-------------|-------------|-----------|------------|
| | | | | |

+ Flow rate values are approximate and are determined using deionized water at 100 cm head

height.

** Pressure injection flow rates are determined using media of 11.8 centipoise (cP) viscosity, with 152 cm administration set tubing.

LBL061293 R01 (2024-05)



Erg[®]Pack. Complete