





(IPNXXXXXX)

## Pressure Injectable One-Lumen PICC



## Contents:

- One Lumen Catheter: 4 Fr. (1.45 mm OD) x 55 cm, Pressure Injectable, T-Port Connector, Blue FlexTip®, Contamination Guard and Placement Wire
- GlideThru™ Peel-Away Sheath: 4 Fr. x 2-3/4" (7 cm) Radiopaque over 4 Fr.
- Spring-Wire Guide: .018" (0.46 mm) dia. x 13-1/8" (33 cm) (Straight Soft Tip on Both Ends)
- Catheter: 22 Ga. x 1-3/4" (4.45 cm) Radiopaque over 25 Ga. RW Introducer
- Introducer Needle: 21 Ga. x 1-1/2" (3.81 cm) TW
- Injection Needle: Eclipse™1 22 Ga. x 1-1/2" (3.81 cm) RW
- Introducer Needle: Echogenic 21 Ga. x 2-3/4" (7 cm) TW
- Filter Needle: 18 Ga. TW x 1-1/2" (3.81 cm) with Blunt Fill Tip
- Injection Needle: Eclipse™1 25 Ga. x 1" (2.54 cm) RW and 3 mL Luer-Lock
- Syringe: 10 mL Luer-Lock
- Syringe: 10 mL Luer-Slip
- Dust Cap: Non-Vented
- SecondSite™ Adjustable Hub: Catheter Clamp
- SecondSite™ Adjustable Hub: Fastener
- SharpsAway® II Locking Disposal Cup
- SharpsAway® Disposal Cup
- Catheter Trimmer
- Towel: 24" x 36" (61 cm x 91 cm)

Not made with natural rubber latex

- Maximal Barrier Drape™ with two 13 cm (5") x 15 cm (6") fenestrations
- Safety Scalpel: #11

contraindications exist.

1: Patient ID Card

1: Sterile Procedure

- Sian
- Patient Information Booklet
- Chart Sticker
- PICC Line Insertion Checklist
- Paper Tape Measure
- Tourniquet
- Gauze Pad: 2" x 2" (5 cm x 5 cm)
- Gauze Pad: 4" x 4" (10 cm x 10 cm)
- Dressing: Tegaderm®2 15.5 cm x 10 cm
- Surgical Apparel: Impervious Gown
- Surgical Apparel: Bouffant Cap
- Surgical Apparel: Mask with Eye Shield
- Dressing: STATLOCK®3 PICC Plus Catheter Stabilization Device
- Surgical Apparel: Mask
- Tape: Steri-Strip®2
- HemoHopper® Fluid Receptacle
- <sup>1</sup>A trademark of Becton, Dickinson and Company
- <sup>2</sup>A registered trademark of 3M Company.
- <sup>3</sup>A registered trademark of C. R. Bard, Inc.


All components are CE 2797 unless otherwise noted.



Lumen	Priming Volume* (mL)	Gravity Flow Rate† (mL/hr)	Pump Flow Rate†† (mL/hr)	MAX Pressure Injection Flow Rate** (mL/sec)
Dietal (40 Ca )	0.40	126	407E	1

<sup>\*</sup> Priming volumes are approximate and are done without accessories.





Contraindications: The Pressure Injectable PICC is contraindicated wherever there is presence of device related infections or presence of

thrombosis in the intended insertion vessel or catheter pathway. Clinical assessment of the patient must be completed to ensure no

LBL063369 R02 (2024-03)

<sup>†</sup> Flow rate values are approximate and are determined using deionized water at 100 cm head

<sup>††</sup> Pump flow rates are determined at maximum pump pressure of 10 psig and represent

<sup>\*\*</sup> Pressure injection flow rates are determined at the injector pressure setting of 300 psi maximum using media of 11.8 centipoise viscosity, with 152 cm pressure tubing.