





(IPNXXXXXX)



Arrowg+ard Blue® Two-Lumen Hemodialysis Catheterization for High Volume **Infusions**



Contents:

- Two-Lumen Catheter: 12 Fr. (4.06 mm OD) x 25 cm Radiopaque Polyurethane with Blue FlexTip®, Arrowg+ard® Antimicrobial Surface Treatment, Extension
- Spring-Wire Guide, Nitinol, Marked: .035" (0.89 mm) dia. x 27-1/2" (70 cm) (Straight Soft Tip on One End - "J" Tip on Other) with Arrow® GlideWheel™ Wire Advancer
- Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer
- Injection Needle: SafetyGlide™1 25 Ga. x 1" (2.54 cm)
- Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm)
- Pressure Transduction Probe
- Introducer Needle: Echogenic 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction Syringe
- Injection Needle: SafetyGlide™1 23 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip
- Syringe: 3 mL Luer-Lock
- Tissue Dilator: 10 Fr. (3.5 mm) x 10.2 cm
- Tissue Dilator: 12 Fr. (4.0 mm) x 14 cm
- 10 mL Luer-Lock Pre-Filled Saline Syringe
- 5 mL 1% Lidocaine HCl Solution and Alcohol Prep
- 3 mL Applicator 2% CHG and 70% IPA ChloraPrep®2 One-Step Solution with

- Hi-Lite Orange™ Tint
- SharpsAway® II Locking Disposal Cup
- SharpsAway® Disposal Cup
- Maximal Barrier Drape™ with 4" fenestration
- 1: Towel
- Needle Holder
- Safety Scalpel: #11
- Checklist/CLIP Sheet
- Sterile Procedure Sign
- Medication Label: 1% Lidocaine
- Gauze Pad: 2" x 2" (5 cm x 5 cm)
- Gauze Pad: 4" x 4" (10 cm x 10 cm)
- Surgical Apparel: Impervious Gown
- Dressing: I.V. Advanced 8.5 cm x 11.5 cm
- Dressing: BIOPATCH®3
- Transducer Cover: 14 cm x 147 cm
- Surgical Apparel: Mask with Eye Shield
- Surgical Apparel: Bouffant Cap
- Suture: 3-0 Silk with Curved Needle
- Tubing±: 20" (50 cm) Extension
- HemoHopper® Fluid Receptacle

- ¹A trademark of Becton, Dickinson and Company.
- ²A registered trademark of CareFusion or one of its subsidiaries.
- ³A registered trademark of Johnson & Johnson Corporation.

±Precaution: Contains Phthalate: DEHP.

Warning: The results of certain animal experiments have shown phthalates to be potentially toxic to reproduction. Proceeding from the present state of scientific knowledge, risks for male premature infants cannot be excluded in the case of long-term exposure or application. Medical products containing phthalates should be used only temporarily with pregnant women, nursing mothers, babies and infants. Not made with natural rubber latex.

Store between 20 - 25°C (68 - 77°F)

Arrow International provides the enclosed medication label(s) for your convenience. Please ensure that these labels are applied to the correct syringe and


Contraindications: The Arrow large-bore two-lumen catheter is not designed for long-term (≥ 30 days) hemodialysis or for use in patients with thrombosed vessels

The Arrowg+ard Blue antimicrobial catheter is contraindicated for patients with known hypersensitivity to chlorhexidine, silver sulfadiazine, and/ or sulfa drugs.



Priming Volume* Distal 1.40 mL Proximal 1.60 mL

* Priming volumes are approximate and are done without accessories.



† In vitro tests performed using 3:1 glycerine/saline at 37°C.















LBL068966 R03 (2024-12)