



7 Fr. **3** Lumen **20** cm catheter length **.032** inch dia. spring-wire guide

Contents:

- | | |
|--|--|
| 1: Three-Lumen Catheter: 7 Fr. (2.3 mm OD) x 20 cm, Pressure Injectable | 1: SharpsAway® II Locking Disposal Cup |
| 1: Spring-Wire Guide, Marked: .032" (0.81 mm) dia. x 23-5/8" (60 cm) (Straight Soft Tip on One End - "J" Tip on Other) with Arrow Advancer | 1: Drape: 41" x 55" with 4" fenestration |
| 1: Catheter: 18 Ga. x 2-1/2" (6.35 cm) Radiopaque over 20 Ga. RW Introducer Needle | 1: Needle Holder |
| 1: Injection Needle: SafetyGlide™ 1 25 Ga. x 1" (2.54 cm) | 1: Safety Scalpel: #11 |
| 1: Blunt Fill Needle: 18 Ga. x 1-1/2" (3.81 cm) | 1: Flow Rate Information Card with Injection Log |
| 1: Pressure Transduction Probe | 1: Medication Label: 1% Lidocaine |
| 1: Introducer Needle: Echogenic 18 Ga. x 2-1/2" (6.35 cm) XTW and 5 mL Arrow® Raulerson Spring-Wire Introduction Syringe | 2: Gauze Pad: 2" x 2" (5 cm x 5 cm) |
| 1: Injection Needle: SafetyGlide™ 1 22 Ga. x 1-1/2" (3.81 cm) and 5 mL Luer-Slip Syringe | 5: Gauze Pad: 4" x 4" (10 cm x 10 cm) |
| 1: Syringe: 3 mL Luer-Lock | 1: Suture: 3-0 Silk with Curved Needle |
| 1: Tissue Dilator: 8.5 Fr. (2.8 mm) x 10.2 cm | 1: Tubing±: 20" (50 cm) Extension |
| 1: 5 mL 1% Lidocaine HCl Solution and Alcohol Prep | |
| 1: 3 mL Applicator 2% CHG and 70% IPA Chloraprep® ² One-Step Solution with Hi-Lite Orange™ Tint | |
| 1: SecondSite™ Adjustable Hub: Fastener | |
| 1: SecondSite™ Adjustable Hub: Catheter Clamp | |
| 1: SharpsAway® Disposal Cup | |

Warning: Read all package insert warnings, precautions, and instructions prior to use. Failure to do so may result in severe patient injury or death.

California Prop. 65
⚠ WARNING: Cancer and Reproductive Harm.
www.P65Warnings.ca.gov


Warning: The results of certain animal experiments have shown phthalates to be potentially toxic to reproduction. Proceeding from the present state of scientific knowledge, risks for male premature infants cannot be excluded in the case of long-term exposure or application. Medical products containing phthalates should be used only temporarily with pregnant women, nursing mothers, babies and infants.

Store between 20 – 25°C (68 – 77°F).

Sterilized using ethylene oxide unless otherwise indicated in the contents list.

Fluid path components are non-pyrogenic.

Arrow International provides the enclosed medication label(s) for your convenience. Please ensure that these labels are applied to the correct syringe and corresponding medication.

 Lumen	Priming Volume* (mL)	Gravity Flow Rate† (mL/hr)	MAX Pressure Injection Flow Rate** (mL/sec)
Distal (16 Ga.)	0.47	2656	10
Medial (18 Ga.)	0.40	1245	5
Proximal (18 Ga.)	0.42	1328	5


† Flow rate values are approximate and are determined using deionized water at 100 cm head height.

** Pressure injection flow rates are determined using media of 11.8 centipoise (cP) viscosity, with 152 cm administration set tubing.



LOT



 Arrow International LLC
Subsidiary of Teleflex Incorporated
3015 Carrington Mill Blvd.
Morrisville, NC 27560 USA

Teleflex®
www.teleflex.com

LBI 069459 R00 (2022-12)



7 Fr. **3** Lumen **20** cm catheter length **.032** inch dia. spring-wire guide

ARROW